




I have no actual or potential conflict of interest to this presentation.

I have nothing to disclose.











	Answer the questions as follows:	
	0 Never	
	1 Seldom	
	2 Sometimes	
	3 Often	
	4 Always	
0		0


0		(
	1. Works constantly, takes little or no time for family and friends, and/or when	
	home, isn't engaged with family members. Seems isolated. Disinterested in	
	social engagement and/or interaction.	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	2. Problems in personal relationships, serious issues with coworkers, feels	
	misunderstood and/or underappreciated.	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	3. Lack of compassion and patience for coworkers, patients and family	
	members, overreaction to minor frustrations, prone to angry outbursts,	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	4. Dreads going to work, discusses work in negative terms or not at all, expresses	
	frustration over workload and/or health records, self-isolating from co-workers,	
	lack of enthusiasm for new patients and/or procedures,	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	Source: AMA Alliance Guide to Physician Burnout and How to Effect Change / www.amaalliance.org/physician-burnout	


9		0
	5. Loss of self-esteem, questions training and skill, disproportionately affected by patients' suffering or loss of life, frustrated about lack of control over professional and/or personal issues, feels trapped. 0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	6. Chronic psychological, emotional or physical fatigue, insomnia, extreme exhaustion at the end of a day, lack of physical and/or emotional energy, increased stress level. 0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	7. Lack of concentration and/or attentiveness. 0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	Increased or new use of alcohol and/or drugs, and/or decreased use of healthy coping mechanisms.	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
0	$Source: AMA\ Alliance\ Guide\ to\ Physician\ Burnout\ and\ How\ to\ Effect\ Change\ /\ {\bf www.amaalliance.org/physician-burnout}$	(6)


9		9
	No longer engages in activities that increase health and wellbeing, such as working out. walking, outdoor activities, relaxation, etc.	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	 Talk of giving up medicine and/or retiring early, wishing he or she had chosen another career, concern about not making a difference, increased cynicism. 	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	11. No longer looks forward to planning and/or taking vacations, time off and/or outside- of-medicine activities.	
	0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
	12. Loss of interest in family, financial, and/or retirement planning. 0 - Never 1 - Seldom 2 - Sometimes 3 - Often 4 - Always	
8	Source: AMA Alliance Guide to Physician Burnout and How to Effect Change / www.amaalliance.org/physician-burnout	
2		9

















It is a sobering fact that an estimated 300 to 400 physicians in the United States commit suicide annually.

Women physicians are at highest risk, with an estimated relative risk ratio of 2.7 for suicide in relation to the general female population.

Schemhammer, et al, Sticide rates among physicians: a quantitative and gender assessment (meta-analysis). AMJ Psychiatry 2004 Dec; 161 (12)



